

The drugs listed below have undesirable side effects and/or may affect your anesthesia or surgery. These drugs should be avoided for two weeks prior to surgery, and two weeks after surgery. If you need to take something for minor aches and pains, you may take Tylenol or another acetaminophen product. **NO ASPIRIN/ASPIRIN-BASED PRODUCTS OR BIRTH CONTROL PILLS/ INJECTIONS\* TWO WEEKS BEFORE AND AFTER.** Aspirin will cause excessive bleeding. Birth control will increase chances of blood clots. **ALSO, TWO WEEKS PRIOR, NO \*\*HERBAL SUPPLEMENTS, VITAMIN E or RED WINE. TESTOSTERONE MUST BE DISCONTINUED 2 WKS BEFORE, BUT MAY BE RESUMED AFTER ONE WEEK.** Please let Dr. Bartlett know if you are currently taking any of these medications on a regular basis.

Advil	Butazolidin	Fiorinal	Pamelor
Aleve	Cama Arthritis Reliever	Fish Oil	Pepto Bismol
Alka Seltzer	Cheracol Capsules	Flagyl	Percodan
Anacin	Chlortrimeton	Flexeril	Perocan
Anaprox	Clinoril	Four Way Cold Tabs	Persistin
Anadynos	Congesprin Chewables	Gaysal S	Phentermine
Ansaid	Cosprin	Ibuprofen	Phenylbutazone
A.P.C.	CP-2 Tabs	Imitrex	Ponseil
Argesic	Damason P	Indocin	Progesterone
Arthropan liquid	Daryon Compound	Indomethacin	Propoxyphene
Arthritis Pain Formula	Darvon	Lanorinal	Compound 45
Arthritis Strength	Darvon N, with A.S.A	Lioresal	Rufen
Bufferin	Dulvules	Lortab	Ru-Tuss
A.S.A.	Di-gesic	Magan	Sine-Aid
A.S.A. Enseals	Disalcid	Magsal	Sinutab
Ascriptin	Dolobid	Measurin	SK-55 Compound
Asperbuf	Doprn	Micrainin	Stendin
Aspergum	Dristan	Midol	Tagamet
Aspirin	Durasal Tablets	Mobigestic	Tamoxifen
Altromid	Easprin	Morbidin	Testosterone
Axotal	Ecotrin	Motrin	Tetracycline
Azolid	Efficin	Mysteelin	Tolectin
Bayer Aspirin	Elavil	Nalfon	Triaminicin
B.C. Tablets and powder	Emagrin	Naprosyn	Trigestic
Birth Control Pills	Empirin with Codeine	Naproxen	Uracel
Birth Control Injection	Emprazil	Neocylate	Vanquish
Buff A Comp	Encaprin	Nicobid	Verin
Bufferin Arthritis	Endep	Norgesic	Vibramycin
Bufferin	Equagesic Tablets	Norgesic Forte	Vitamin E
Bufferin Extra Strength	Estrogen Blockers	Nuprin	Voltaren
Buffet II	Etrafon	Oraflex	Zomax
Buf-Tabs	Excedrin	Orudis	Zorprin
	Feldene	Osteo Bi-Flex	

***Please use this list as a guideline. Report the use of all medications to Dr. Bartlett 617-735-1800.***

\*Birth control injections cannot be administered within two weeks of surgery (2 weeks before and after).

\*\*Herbal supplements are not regulated by the FDA, therefore we do not know for certain what the supplements contain. In order to ensure patient safety, we advise stopping all herbal supplements two weeks prior to your surgical procedure.